



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## 266 *Publications of the Astronomical Society, &c.*

operating the hand clamps and shifting certain levers connected with the gear train in a certain specified order. An automatic electrically operated gear-shifting mechanism has been designed, and is in process of construction, which will make these changes at the proper time, with no attention from the observer.

H. D. CURTIS.

### CORRECTION.

In my note on the unit of stellar distance in the August number of the *Publications* I regret that I inadvertently grouped the term *siriometer* with other terms which have been suggested as names for the distance corresponding to a parallax of one second of arc. The name *siriometer* was introduced by Professor C. V. L. CHARLIER, and is the distance equal to one million times the mean distance of the Sun from the Earth. It thus is almost identical with the term *Siriusweite* (the distance corresponding to a parallax of 0.2 seconds of arc); the parallax of a star at a distance of one *siriometer* would be equal to  $0''.206 \pm$ . It has the advantage that it is a definite and easily remembered multiple of the astronomical unit.

H. D. CURTIS.

### REDISCOVERY OF WESTPHAL'S PERIODIC COMET.

The fourth comet of the year was discovered on September 26, 1913, by Mr. DELAVAN, at the La Plata Observatory, Argentina. This proves to be identical with WESTPHAL's comet of 1852, which was expected to return to the neighborhood of the Earth this year, as the computed period was about sixty-one years. Only three comets of longer period than this have so far been observed at a second return to perihelion. These are HALLEY's comet, PONS-BROOKS' comet (1812-1883), and OLBERS-BROOKS' comet (1815-1887). In 1852, WESTPHAL's comet was, for a time, visible to the naked eye, though not a great comet.

R. G. AITKEN.